

Solution of matrix models
in
the DIII generator ensemble

Harold Roussel¹

Physics Department, McGill University
Ernest Rutherford Building
Montréal, Québec H3A 2T8
Canada

Abstract

In this work we solve two new matrix models, using standard and new techniques. The two models are based on matrix ensembles not previously considered, namely the DIII generator ensemble. It is shown that, in the double scaling limit, their free energy has the same behavior as previous models describing oriented and unoriented surfaces. We also found an additional solution for one of the models.

¹roussel@hep.physics.mcgill.ca

Introduction

Recent progress has been made in non-perturbative string theory via matrix models, where one represents triangulated random surfaces by large- N matrix integrals. By taking the so-called double scaling limit, one gets some insight in the physics of the model, associated with models of 2-D gravity. In this paper we look at two new such matrix models.

The reasons for studying them are that they were part of a classification scheme, but also they provided an opportunity to extend the techniques for solving matrix models, namely via skew-orthogonal polynomials [1]. We also had to solve them to compare with previous models to see if the physics revealed by the free energy is the same. Indeed, from its series expansion, we can say if the model describes unoriented surfaces in addition to oriented ones, and, by calculating an appropriate ratio, we can say if it is the same as previous models.

Section one is devoted to a classification scheme for matrix models within the context of symmetric spaces. In section two and three, we show the calculations of two new models in $D = 0$. Finally, we conclude with a discussion and a comparison of the two models' free energies, and a comparison with other models.

1 Classification of matrix models

We will now introduce a classification of matrix models based on symmetric spaces. This is an interesting class of single-matrix models for which the reduction of the matrix integral allows for a solution by a polynomial method.

Symmetric spaces are cosets of the form G/K . Technically speaking, those that we are considering are the simply connected Riemannian globally symmetric spaces [2, 3, 11]. The classical types are listed in table 1. The two matrix models, in sections two and three, are based on the DIII generator ensemble. They correspond, for section two, to n odd, and for section three, to n even.

There are a few ways to reduce a matrix integral, depending on the type of matrix or the matrix ensemble. For example, for the Hermitian matrix model, one can simply diagonalize the matrix as it is done in Metha's book [4]. The result is that we get an integral over the eigenvalues with a Jacobian in terms of these eigenvalues. The idea behind this reduction is to simplify the matrix integral.

The symmetric spaces introduced in the previous subsection give us a systematic

approach to identify the matrix ensemble for which a similar reduction is possible. For these spaces, however, the procedure of reducing the matrix integral is more complex. An analogy can be used to understand how the Jacobian and the set of integration parameters arise. Consider the volume element in flat space written in cartesian coordinates and in spherical coordinates, $\prod_{i=1}^N dx_i = d\omega dr r^{N-1}$. Spherical symmetry would allow us to integrate out the angular part $d\omega$ with r^{N-1} being the Jacobian and r the integration variable. For a symmetric space (matrix ensemble) the Haar measure can be decomposed in the same way, with the symmetry provided by the associated group. In doing so we get the Jacobian and also a set of parameters (which are not necessarily the eigenvalues of the matrices). The example above can be regarded as the measure for the generator ensemble of the coset $SO(N+1)/SO(N)$. The parameter of interest is r and $d\omega$ is the measure on $SO(N)$, which yields a “trivial” integral if the original integrand is spherically symmetric. Such vector models were studied in [5]. For the technical details the reader is referred to Helgason’s book [3]. Let us just say here that there is a relation between the multiplicities of the roots and the exponents in the Jacobian. Table 2 gives some symmetric spaces and their associated Jacobian. Many of these models have been previously considered, including A I [6, 7], corresponding to real symmetric matrices, A II [7, 8], corresponding to quaternionic real self dual matrices, as well as A III, BD I, and C II [5, 9].

It should be noted here that matrix models with Hermitian, antisymmetric, and quaternionic real self dual matrices, are not part of this classification. In fact, they are generator ensembles for the classical groups $U(n)$, $SO(n)$, $Sp(2n)$, respectively.

In the present paper we consider the D III generator ensemble, so only C I remains to be studied.

2 First model – n odd

We will first study the double-scaling limit of the D III generator ensemble with n odd (see table 2). The matrix integral always takes the form,

$$\mathcal{Z} = \int dM e^{-\beta \text{Tr} V(M)}. \quad (1)$$

The Jacobian for this model is

$$\mathcal{J} = \prod_i^N |x_i^5| \prod_{i < j}^N (x_i^2 - x_j^2)^4, \quad (2)$$

and we are going to make the analysis with the potential $V = ax^2 + b/2 x^4$. The partition function can then be written as follows,

$$\mathcal{Z}_N = 2^{-N} \int_0^\infty \prod_i^N dy_i \prod_i^N y_i^2 \prod_{i<j}^N (y_i - y_j)^4 e^{-\beta(ay_i + b/2 y_i^2)}, \quad (3)$$

where we made the change of variables $x^2 = y$. For convenience of evaluation we will rewrite it as a determinant [4]. We have,

$$\prod_i^N y_i^2 \prod_{i<j}^N (y_i - y_j)^4 = \det \begin{bmatrix} P_0(y_1) & P_1(y_1) & \dots & P_{2N-1}(y_1) \\ y^2 \partial_y P_0(y_1) & y^2 \partial_y P_1(y_1) & \dots & y^2 \partial_y P_{2N-1}(y_1) \\ P_0(y_2) & P_1(y_2) & \dots & P_{2N-1}(y_2) \\ \vdots & \vdots & & \vdots \\ y^2 \partial_y P_0(y_N) & y^2 \partial_y P_1(y_N) & \dots & y^2 \partial_y P_{2N-1}(y_N) \end{bmatrix}. \quad (4)$$

The P 's are the usual orthogonal polynomials on the half-line, defined as

$$P_k(y) = y^k + l.o. \quad P_0(y) = 1 \quad (5)$$

$$\int_0^\infty d\mu P_k(y) P_l(y) = h_k \delta_{l,k} \quad d\mu = dy e^{-\beta V(y)} \quad (6)$$

and from which we derive the following recursion relation,

$$y P_k = P_{k+1} + S_k P_k + R_k P_{k-1}, \quad (7)$$

with $R_k = h_k/h_{k-1}$.

In order to solve for the free energy, we will need to know, first, the solutions for R and S . By considering recursion relations for $\partial_y P_k$ and $y \partial_y P_k$, one finds a set of two self-consistent equations in terms of R , S , k , a , b , and β [10],

$$\frac{4k+1}{\beta} = a S_{2k} + b(R_{2k} + R_{2k+1} + S_{2k}^2) \quad (8)$$

and

$$(a + b S_{2k})(a + b S_{2k-1}) R_{2k} = \frac{1}{4} [S_{2k}(a + b S_{2k}) + b(R_{2k+1} - R_{2k}) - \frac{1}{\beta}]^2. \quad (9)$$

We used $2k$ indices because in the final recursion relation, we will need to know the solutions of R_{2N+l} and S_{2N+l} . In the planar approximation, and choosing $\lambda_c = (k/\beta)_c = 1/4$ and $R_c = 1$ (together with the criticality condition $\partial_S \lambda = 0$), we find all the critical values,

$$\begin{aligned} \lambda_c &= 1/4 & R_c &= 1 & S_c &= 2 \\ a &= 1 & b &= -1/6. \end{aligned} \quad (10)$$

We choose the scaling solutions to be,

$$S_{2N+l} = 2(1 - \beta^{-\mu} \exp(-\frac{l}{2}\beta^{-\nu} \frac{\partial}{\partial t})f) \quad (11)$$

$$R_{2N+l} = 1 - \beta^{-\mu} \exp(-\frac{l}{2}\beta^{-\nu} \frac{\partial}{\partial t})(g_0 + \beta^{-\nu} g_1 + g_2 \beta^{-2\nu} + \dots) \quad (12)$$

where t is defined by $t = (\beta/4 - N)\beta^{-\nu}$. We write N instead of k because we are only interested in the large N behavior of R and S (and so we replace k by N in our recursion relations as well). In the R ansatz we added some more degrees of freedom by expanding the g function in powers of $\beta^{-\nu}$. This will be necessary to get consistent solutions beyond leading order in $\beta^{-\nu}$. Doing the same thing for f would only yield redundant equations.

After inserting in the recursion relations, and working out the lowest order equations, we see that a consistent solution requires $\mu = 2\nu$. For the two recursion relations, the results are given, order by order, in table 3.

From eq.(8), we obtain the values of the exponents, $\nu = 1/5$, and $\mu = 2/5$, which is consistent with our previous relation between μ and ν . Also, using the relations between the g coefficients and f , we get, at order $\beta^{-\nu-2\nu}$ in (9), a differential equation for f ,

$$4t = f^2 - 1/48 f'' \quad (13)$$

which is the well-known Painlevé I equation, upon renormalization of t . At order $\beta^{-\mu}$ and $\beta^{-\mu-\nu}$, we simply get equalities (e.g. $0 = 0$), which come from our criticality requirement (first derivative of λ with respect to S is zero).

To solve the model, one must still solve for the double-scaling limit of the entire partition function. For this, it is clear that we have to find a recursion relation for $y^2 \partial_y P_n(y)$ (as appears in the determinant 4). After some algebraic manipulations, one finds

$$\begin{aligned} y^2 \partial_y P_n = & n P_{n+1} + \left(\frac{\beta}{2} C_{n,n} - S_n \right) P_n + (\beta C_{n-1,n} - n - 1) R_n P_{n-1} + \\ & \beta C_{n-2,n} R_n R_{n-1} P_{n-2} + \beta C_{n-3,n} R_n R_{n-1} R_{n-2} P_{n-3}, \end{aligned} \quad (14)$$

where, for simplicity we wrote $C_{k,l} = \int d\mu P_k P_l y^2 V'(y)$. With this recursion relation, the partition function is found to be,

$$\begin{aligned} \mathcal{Z}_{N+1} = & (N+1)(\langle P'_{2N}, P_{2N+1} \rangle - \langle P'_{2N+1}, P_{2N} \rangle) \mathcal{Z}_N + N(N+1) \\ & \langle P'_{2N-2}, P_{2N+1} \rangle (\langle P'_{2N-1}, P_{2N} \rangle - \langle P'_{2N}, P_{2N-1} \rangle) \mathcal{Z}_{N-1} \\ & - N(N+1)(N-1) \langle P'_{2N-2}, P_{2N+1} \rangle \langle P'_{2N-3}, P_{2N} \rangle \\ & \langle P'_{2N-4}, P_{2N-1} \rangle \mathcal{Z}_{N-2} - (N+1) \langle P'_{2N-1}, P_{2N+1} \rangle Y_N, \end{aligned} \quad (15)$$

where $P' = y^2 \partial_y P$. Y_N is an auxilliary partition function with a determinant similar to Z_N but where the last and third-to-last columns were removed. It was introduced to avoid an infinite number of terms in the previous relation, and satisfies

$$Y_N = N < P'_{2N-2}, P_{2N} > \mathcal{Z}_{N-1} - N < P'_{2N-3}, P_{2N} > Y_{N-1}. \quad (16)$$

We now define, for later convenience (we want to have a smooth planar limit as $N \rightarrow \infty$), the following two ansatz,

$$W_N = \frac{\mathcal{Z}_N}{\mathcal{Z}_{N-1} N \beta b h_{2N-1}} \quad X_N = \frac{Y_N}{\mathcal{Z}_{N-1} N \beta b h_{2N-1}}, \quad (17)$$

and rewrite eqs.(15) and (16) in terms of polynomials in W and X . After some algebraic manipulations, and expanding the brackets, we get

$$\begin{aligned} W_{N+1} W_N W_{N-1} - \frac{W_N W_{N-1}}{b} (a(S_{2N+1} + S_{2N}) + b(R_{2N} + R_{2N+1} + R_{2N+2} + S_{2N+1}^2 \\ + S_{2N}^2 + S_{2N+1} S_{2N}) - \frac{4N}{\beta} - \frac{2}{\beta}) + \frac{X_N W_{N-1}}{b} (a + b(S_{2N+1} + S_{2N} + S_{2N-1})) \\ - W_{N-1} R_{2N} (R_{2N+1} + R_{2N} + R_{2N-1} + S_{2N-1}^2 + S_{2N}^2 + S_{2N-1} S_{2N}) \\ - \frac{W_{N-1} R_{2N}}{b} \left(a(S_{2N} + S_{2N-1}) - \frac{4N}{\beta} \right) + R_{2N} R_{2N-1} R_{2N-2} = 0 \end{aligned} \quad (18)$$

$$X_{N+1} W_N - \frac{R_{2N+2} W_N}{b} (a + b(S_{2N+2} + S_{2N+1} + S_{2N})) + R_{2N+2} X_N = 0. \quad (19)$$

In the planar limit, and using the previously found critical values for R , S , etc, we find that the critical values are $W_c = -1$ (triple root of 18), while X_c remains undetermined. X it will be determined below when solving the recursion relations.

To solve them, we choose the following ansatz,

$$W_{N+l} = -1 + \beta^{-\rho} \exp(-l\beta^{-\nu} \frac{\partial}{\partial t})(h_0 + \beta^{-\nu} h_1 + \beta^{-2\nu} h_2 + \dots) \quad (20)$$

$$X_{N+l} = X_c - \beta^{-\sigma} \exp(-l\beta^{-\nu} \frac{\partial}{\partial t})(k_0 + \beta^{-\nu} k_1 + \beta^{-2\nu} k_2 + \dots). \quad (21)$$

After insertion in eqs. (18) and (19) (at $\beta^{-2\nu}$ order), and using $\phi = \mu = 2/5$, $\sigma = \rho = \nu = 1/5$, we find the results given in table 4. We see that a consistent solution requires that $X_c = 0$. In doing so, we get two differential equations defining h_0 and k_0 ,

$$0 = 6f + h_0 k_0 + k_0' \quad (22)$$

$$0 = -6f' - 12f h_0 + h_0^3 + 3h_0 h_0' + h_0'' - 6f k_0. \quad (23)$$

We used the fact that $g_0 = 2f$ (table 3). Using the known solution for f ,

$$f = 2t^{1/2} - \frac{1}{384}t^{-2} - \frac{49}{589824}t^{-9/2} - \dots, \quad (24)$$

and power series solution for h_0 and k_0 , we find a set of algebraic equations that we can solve. We finally end up with four solutions,

$$\begin{aligned} h_0 &= \pm 2\sqrt{3}t^{1/4} + \frac{7}{8}t^{-1} \mp \frac{5\sqrt{3}}{384}t^{-9/4} + \dots \\ k_0 &= \mp 2\sqrt{3}t^{1/4} + \frac{9}{8}t^{-1} \pm \frac{5\sqrt{3}}{384}t^{-9/4} + \dots \end{aligned} \quad (25)$$

or

$$\begin{aligned} h_0 &= \pm 2\sqrt{3}t^{1/4} - \frac{1}{8}t^{-1} \mp \frac{5\sqrt{3}}{384}t^{-9/4} + \dots \\ k_0 &= \mp 2\sqrt{3}t^{1/4} + \frac{1}{8}t^{-1} \pm \frac{5\sqrt{3}}{384}t^{-9/4} + \dots \end{aligned} \quad (26)$$

It turns out that we will only need h_0 in the solution of the free energy. Using the following relation for large N ,

$$\exp(-\partial_N^2 F) = \frac{\mathcal{Z}_{N+1}\mathcal{Z}_{N-1}}{\mathcal{Z}_N^2} = R_{2N+1}R_{2N}\frac{W_{N+1}}{W_N}, \quad (27)$$

the second derivative of the free energy is found to be

$$F'' \simeq 4f - h'_0. \quad (28)$$

Using the solution for f and the solution (26) for h_0 , we find,

$$F'' = 8t^{1/2} \mp \frac{\sqrt{3}}{2}t^{-3/4} - \frac{13}{96}t^{-2} + \dots \quad (29)$$

In order to compare this result with other matrix models, we need a quantity independent of the scale of t and F'' . Such a quantity is the product of the first and the third coefficient in the series for F'' divided by the square of the second coefficient. Here we find $-13/9$, which is the same result as in [6, 7] although the matrix ensemble in these papers is different. From previous solutions this is the expected ratio for the free energy of pure 2D quantum gravity with oriented and unoriented surfaces. Doing the same with (25), we get,

$$F'' = 8t^{1/2} \mp \frac{\sqrt{3}}{2}t^{-3/4} + \frac{83}{96}t^{-2} + \dots \quad (30)$$

In that case, the universal ratio c_0c_2/c_1^2 yields $83/9$, which differs from any known matrix models. One surprising result is that our two solutions differ only by t^{-2} (verified up to 15th order of F''), which is the term corresponding to the torus and the Klein bottle.

3 Second model – n even

We start again from the matrix integral (1). But now, the Jacobian takes the form,

$$\mathcal{J} = \prod_i^N |x_i| \prod_{i<j}^N (x_i^2 - x_j^2)^4. \quad (31)$$

We are doing the analysis, as usual, with the potential $V = ax^2 + b/2 x^4$. The partition function can then be written as follows,

$$\mathcal{Z}_N = 2^{-N} \int_0^\infty \prod_i^N dy_i \prod_{i<j}^N (y_i - y_j)^4 e^{-\beta(ay_i + b/2 y_i^2)}. \quad (32)$$

We used, again, the substitution $x^2 = y$. The complicated part is the Jacobian. For convenience of evaluation we rewrite it as a determinant [1]. We have,

$$\prod_{i<j}^N (y_i - y_j)^4 = \det \begin{bmatrix} Q_0(y_1) & Q_1(y_1) & \dots & Q_{2N-1}(y_1) \\ \partial_y Q_0(y_1) & \partial_y Q_1(y_1) & \dots & \partial_y Q_{2N-1}(y_1) \\ Q_0(y_2) & Q_1(y_2) & \dots & Q_{2N-1}(y_2) \\ \vdots & \vdots & & \vdots \\ \partial_y Q_0(y_N) & \partial_y Q_1(y_N) & \dots & \partial_y Q_{2N-1}(y_N) \end{bmatrix}, \quad (33)$$

where the Q 's are Metha's skew-orthogonal polynomials,

$$Q_i(y) = y^i + l.o., \quad (34)$$

and

$$\langle Q_i, Q_j \rangle_Q \equiv \frac{1}{2} \int dy e^{-\beta V(y)} (Q_i Q'_j - Q'_i Q_j) = q_{[i/2]z_{ij}}, \quad (35)$$

with

$$z_{2i,2i+1} = 1 \quad (36)$$

$$z_{2i+1,2i} = -1, \quad (37)$$

all others being 0. With these definitions we can easily evaluate $\mathcal{Z}_N$,

$$\mathcal{Z}_N = 2^{-N} N! \prod_{i=0}^{N-1} q_i. \quad (38)$$

Here we used skew-orthogonal polynomials because with the orthogonal ones, $\mathcal{Z}_N$ cannot be evaluated (i.e. gives an infinite recursion relation for ∂P). On the other hand,

with Q polynomials, $\mathcal{Z}_N$ is easy to find, but the problem is to establish recursion relations for them and the q 's. Indeed, the only known recursion relation is infinite [1]. The approach that we will follow is to relate the Q polynomials with the P polynomials (these are the usual orthogonal polynomials for which recursion relations are well-known, or at least, easy to find).

We start with the general expansion of the P 's in term of the Q 's,

$$P_{2i} = Q_{2i} + \omega_{i1}Q_{2i-1} + \omega_{i2}Q_{2i-2} + \dots \quad (39)$$

$$P_{2i+1} = Q_{2i+1} + \xi_{i1}Q_{2i} + \xi_{i2}Q_{2i-1} + \dots \quad (40)$$

In order to find odd and even ξ 's and ω 's, we consider four Q products. In the calculation we take the Q product of a P polynomial and a Q polynomial. After using the expansion of P 's in terms of Q 's, and rewriting the Q product in a P product (orthogonal polynomial relation), we get the following relations,

$$\xi_{i,2j}q_{i-j} = \frac{1}{2} \langle \beta V' Q_{2i-2j}, P_{2i+1} \rangle_P - \frac{1}{2} Q_{2i-2j}(0) P_{2i+1}(0) \quad (41)$$

$$\omega_{i,2j-1}q_{i-j} = \frac{1}{2} \langle \beta V' Q_{2i-2j}, P_{2i} \rangle_P - \frac{1}{2} Q_{2i-2j}(0) P_{2i}(0) \quad (42)$$

$$\begin{aligned} -\omega_{i,2j}q_{i-j} &= \frac{1}{2} \langle \beta V' Q_{2i-2j+1}, P_{2i} \rangle_P - \frac{1}{2} Q_{2i-2j+1}(0) P_{2i}(0) \\ &\quad - \frac{1}{2} (2i+1) \delta_{j0} h_{2i} \end{aligned} \quad (43)$$

$$-\xi_{i,2j+1}q_{i-j} = \frac{1}{2} \langle \beta V' Q_{2i-2j+1}, P_{2i+1} \rangle_P - \frac{1}{2} Q_{2i-2j+1}(0) P_{2i+1}(0). \quad (44)$$

Considering the above equations for specific values of j , and defining the following quantities which have a smooth planar limit,

$$\begin{aligned} W_i &= \frac{q_i}{\beta h_{2i}} & X_i &= \frac{Q_{2i}(0)}{P_{2i}(0)} & Y_i &= \frac{Q_{2i+1}(0)}{P_{2i}(0)} \\ & & Z_k &= \frac{P_{k+1}(0)}{P_k(0)} & A_k &= \frac{P_k(0)^2}{\beta h_k}, \end{aligned} \quad (45)$$

we finally get,

$$\frac{W_i}{R_{2i+1}} = \frac{b}{2} - \frac{1}{2} \frac{X_i A_{2i+1}}{Z_{2i}} \quad (46)$$

$$\frac{\xi_{i2} W_{i-1}}{R_{2i+1} R_{2i} R_{2i-1}} = -\frac{1}{2} \frac{X_{i-1} A_{2i+1}}{Z_{2i} Z_{2i-1} Z_{2i-2}} \quad (47)$$

$$0 = a + b S_{2i} - \omega_{i1} b - X_i A_{2i} \quad (48)$$

$$\frac{\omega_{i1} W_{i-1}}{R_{2i} R_{2i-1}} = -\frac{1}{2} \frac{X_{i-1} A_{2i}}{Z_{2i-1} Z_{2i-2}} \quad (49)$$

$$-2W_i = bR_{2i+1} - \xi_{i1}(a + bS_{2i}) + (\xi_{i1}\omega_{i1} - \xi_{i2})b - Y_i A_{2i} - \frac{(2i+1)}{\beta} \quad (50)$$

$$\frac{-\omega_{i2}W_{i-1}}{R_{2i}R_{2i-1}} = \frac{b}{2} - \frac{1}{2} \frac{Y_{i-1}A_{2i}}{Z_{2i-1}Z_{2i-2}} \quad (51)$$

$$\frac{-2\xi_{i1}W_i}{R_{2i+1}} = a + bS_{2i+1} - b\xi_{i1} - \frac{Y_i A_{2i+1}}{Z_{2i}}. \quad (52)$$

In the planar limit, and using critical values found in the previous section, we find,

$$\begin{aligned} b &\rightarrow -1/6 & \lambda = i/\beta &\rightarrow 1/4 & S &\rightarrow 2 & Z &\rightarrow -1 & \omega_{i,n} &\rightarrow \omega_n \\ a &\rightarrow 1 & R &\rightarrow 1 & A &\rightarrow 2/3 & \xi_{i,n} &\rightarrow \xi_n \\ X &\rightarrow 1/2 & W &\rightarrow 1/12 & \omega_1 &\rightarrow -2 & \xi_2 &\rightarrow 2 & \xi_1 &\rightarrow -3/2 \\ & & & & Y &\rightarrow 1/2 & \omega_2 &\rightarrow 3, \end{aligned} \quad (53)$$

where the values for A and Z can be easily found using their explicit form.

We can take a look at the partition function to see exactly what quantities we have to know. From (38), and using (45), we get

$$\frac{\mathcal{Z}_{N+1}\mathcal{Z}_{N-1}}{\mathcal{Z}_N^2} = (1 + \frac{1}{N})R_{2N}R_{2N-1}\frac{W_N}{W_{N-1}}. \quad (54)$$

However, the ratio of $\mathcal{Z}$'s is related to F'' , so all we have to know is the ratio W_N/W_{N-1} , something that we can easily find with a recursion relation for the W 's (of course we also have to find the differential equation satisfied by the function used in the W ansatz).

To find this recursion relation we only need eqs.(46), (48), and (49). After some algebraic manipulations we get a recursion relation for W_N ,

$$\begin{aligned} &(-\frac{(2N+1)}{\beta} + bR_{2N+1})(-\frac{2N}{\beta} + bR_{2N})[2W_N - bR_{2N+1}]W_{N-1} \\ &+ R_{2N+1}(a + bS_{2N})(a + bS_{2N+1})[(-\frac{2N}{\beta}W_{N-1}) + \frac{b^2}{2}R_{2N}R_{2N-1}] = 0. \end{aligned} \quad (55)$$

Using some information from table 3, which applied for recursion relations of P 's, and using $\phi = \mu = 2/5$, $\rho = \nu = 1/5$, we find the coefficients listed in table 5. Replacing solutions for R_{2N+l} , and W_{2N+l} in (54), we get,

$$F'' = 4f - h'_0. \quad (56)$$

Using a power series expansion for f and replacing in $6f = h_0^2 + h'_0$, we find

$$h_0 = \pm 2\sqrt{3}t^{1/4} - \frac{1}{8}t^{-1} \mp \frac{5\sqrt{3}}{384}t^{-9/4} + \dots \quad (57)$$

Finally, we end up with the following solution for the second derivative of the free energy,

$$F'' = 4f - h'_0 \simeq 8t^{1/2} \mp \frac{\sqrt{3}}{2}t^{-3/4} - \frac{13}{96}t^{-2} + \dots \quad (58)$$

The appearance of a term $O(t^{-3/4})$ reveals that this model describes unoriented surfaces for sure, as well as oriented ones. To compare with previous model, we consider the ratio of coefficients,

$$\frac{c_0 c_2}{c_1^2} = \frac{8 \times -\frac{13}{96}}{(\mp \frac{\sqrt{3}}{2})^2} = -\frac{13}{9}. \quad (59)$$

This is the same ratio as what was found in the first solution of the previous section so it describes exactly the same physics.

4 Conclusion

We now summarize our two main results, with their implications. Firstly, for each of the models, the free energy (up to overall scalings) was the same as other models previously studied and using completely different matrix ensembles, as revealed by the ratio $\frac{c_0 c_2}{c_1^2} = -13/9$). This means the following: if these models were actually describing gravity coupled to some other system, then we expect that different regulators would introduce a dependence of the free energy on a coupling parameter. Hence using different matrix ensembles would yield different results. But this is not the case so our analysis confirms that all of these models describe pure gravity including both oriented and unoriented surfaces, as it was first assumed for previous solutions.

Secondly, although both models yield exactly the same result, there is an additional solution for n -odd (ratio $83/9$) . This solution differs from the other one only by the coefficient of the torus/Klein bottle term. So all ratios that do not involve this term are the same in both solutions. A similar result was found in [8] for QRSD matrices, where one solution did not include odd Euler character surfaces. In the present case, the physical interpretation of the extra solution remains unclear.

So far, the C I generator ensemble remains unstudied (as are most of the circular ensembles, which integrate over the entire symmetric space [5]). An interesting observation is that measures for D III - n even - and C I are similar to A II and A I respectively after a change of variables $y = x^2$. The D III and C I generator ensembles are more complicated to analyse though, because their range of integration is from 0 to ∞ instead of $-\infty$ to ∞ , which causes the appearance of boundary terms in the recursion relations. Given that D III with n even and A II yield the same physics in the double-scaling limit, we might expect C I and A I to do as well.

Acknowledgment

I would like to thank Robert C. Myers, for his help throughout the completion of this work. This paper is a shortened version of a thesis presented in partial fulfillment for the degree of master in physics. This research was supported by NSERC.

References

- [1] M. L. Metha and G. Mahoux. *Indian journal of pure and applied math.* **22(7)**, 531, (1990).
- [2] M. A. Olshanetsky and A. M. Perelomov. *Phys. Rep.* **94**, 313–404, (1983).
- [3] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces.* Academic Press, New York, 1978.
- [4] M. L. Metha. *Random matrices.* Academic Press, New York, 1967.
- [5] Arlen Anderson, Robert C. Myers, and Vipul Periwal. *Phys. Lett.* **B254**, 89, (1991).
 Arlen Anderson, Robert C. Myers, and Vipul Periwal. *Nucl. Phys.* **B360**, 463, (1991).
 S. Nishigaki and T. Yonega. *Nucl. Phys.* **B348**, 787, (1991).
 P. Di Vecchia, M. Kato, and N. Ohta. *Nucl. Phys.* **B356**, 495, (1991).
- [6] E. Brézin and H. Neuberger. *Nucl. Phys.* **B350**, 513, (1991).
- [7] E. J. Martinec and G. R. Harris. *Phys. Lett.* **B245**, 384, (1990).
- [8] Robert C. Myers and Vipul Periwal. *Phys. Rev. Lett.* **64**, 3111, (1990).
- [9] Robert C. Myers. *Nucl. Phys.* **B368**, 701, (1992).
 Robert C. Myers and Vipul Periwal. *Nucl. Phys.* **B390**, 716, (1993).
- [10] T.R. Morris. *Nucl. Phys.* **B356**, 703, (1991).
- [11] R. Gilmore. *Lie Groups, Lie Algebras, and Some of Their Applications.* John Wiley and Sons, New York, 1974.

Cartan's notation	$X = G/K$	Rank	System of restricted roots	Multiplicity
A I	$SL(n, R)/SO(n)$	$n - 1$	A_{n-1}	$m_\alpha = 1$
A II	$SU(2n)/Sp(n)$	$n - 1$	A_{n-1}	$m_\alpha = 4$
A III	$SU(p, q)/S(U(p) \times U(q))$	$n=\min(p,q)$	$p = 1, C_n$ $p > q, BC_n$	$m_\alpha = 2, m_{2\beta} = 1$ $m_\alpha = 2, m_{2\beta} = 1$ $m_\beta = 2(p - q)$
BD I	$SO(p, q)/SO(p) \times SO(q)$	$n=\min(p,q)$	$p = q, D_n$ $p > q, B_n$	$m_\alpha = 1$ $m_\beta = (p - q)$ $m_\alpha = 1$
D III	$SO(2n)/U(n)$	$n/2$	$n = 2k, C_k$ $n = 2k + 1$ BC_k	$m_\alpha = 4, m_{2\beta} = 1$ $m_\alpha = 4, m_{2\beta} = 1$ $m_\beta = 4$
C I	$Sp(n, R)/U(n)$	n	C_n	$m_\alpha = 1$
C II	$Sp(p, q)/Sp(p) \times Sp(q)$	$n=\min(p,q)$	$p = q, C_n$ $p > q, BC_n$	$m_\alpha = 4, m_{2\beta} = 3$ $m_\alpha = 4, m_{2\beta} = 3$ $m_\beta = 4(p - q)$

Table 1: Classification of the symmetric spaces.

Cartan's notation	System of restricted roots	Jacobian for generator ensemble
A I	A_{n-1}	$\prod_{i < j} x_i - x_j $
A II	A_{n-1}	$\prod_{i < j} x_i - x_j ^4$
A III	$p = 1, C_n$	$\prod_{i < j} (x_i^2 - x_j^2)^2 \prod x_i ^{2(1-q)+1}$
	$p > q, BC_n$	$\prod_{i < j} (x_i^2 - x_j^2)^2 \prod x_i ^{2(p-q)+1}$
BD I	$p = q, D_n$	$\prod_{i < j} x_i^2 - x_j^2 $
	$p > q, B_n$	$\prod_{i < j} x_i^2 - x_j^2 \prod x_i ^{p-q}$
C I	C_n	$\prod_{i < j} x_i^2 - x_j^2 \prod x_i $
C II	$p = q, C_n$	$\prod_{i < j} (x_i^2 - x_j^2)^4 \prod x_i ^3$
	$p > q, BC_n$	$\prod_{i < j} (x_i^2 - x_j^2)^4 \prod x_i ^{4(p-q)+3}$
D III	$n = 2k, C_k$	$\prod_{i < j} (x_i^2 - x_j^2)^4 \prod x_i $
	$n = 2k + 1, BC_k$	$\prod_{i < j} (x_i^2 - x_j^2)^4 \prod x_i^5 $

Table 2: Symmetric spaces and their Jacobians.

order	eq.(9)	eq.(8)
β^0	$4/9 = 4/9$	$1=1$
$\beta^{-\mu}$	$g_0 = 2f$	$g_0 = 2f$
$\beta^{-\mu-\nu}$	$g_1 = f'/2$	$g'_0 = 4g_1$
$\beta^{-2\mu} = \beta^{-\mu-2\nu}$	$g_2 = 1/16 f'' - f^2$	$4t = 2/3 f^2 - 1/3 g_2 + 1/12 g'_1 - 1/48 g''_0$

Table 3: Lowest order solutions of the two recursion relations for R_{2N+l} and S_{2N+l} .

order	eq.(18)	eq.(19)
$\beta^{-1/5}$	0	$h_0 X_c = 0$
$\beta^{-2/5}$	$6f X_c = 0$	$-6f - g_0 X_c + h_1 X_c - h_0 k_0 - k'_0 = 0$
$\beta^{-3/5}$	$-3g'_0 - 6g_0 h_0 + h_0^3 + 3h_0 h'_0 + h''_0 - 6fk_0 - 6fh_0 X_c = 0$	$3f' + 6fh_0 + g_0 k_0 - h_1 k_0 + h_0 k'_0 - h_0 k_1$ $k''_0/2 - g_1 X_c + g'_0 X_c + h_2 X_c - k'_1 = 0$

Table 4: Lowest order solutions of the two recursion relations for the DIII – odd n – matrix model.

order	eq.(55)
$\beta^{-1/5}$	0
$\beta^{-2/5}$	$6f = h_0^2 + h'_0$

Table 5: Lowest order solutions of the recursion relation for the DIII – even n – matrix model.